

# **Theoretical progress and status on pion scattering length**

**Nicola Cabibbo**


# Summary

- Pion scattering lengths and the cusp.
- Unitarity and analyticity.
- $\pi\pi$  scattering.
- $K \rightarrow 3\pi$  amplitudes at  $O(a_i^2)$ .

N. Cabibbo, *Phys. Rev. Lett.*, 93:121801, 2004.

N. Cabibbo and G. Isidori, *J. High Energy Phys.* JHEP03(2005)021

## How the Cusp looks



The  $\pi^0\pi^0$  spectrum in  $K^+ \rightarrow \pi^+\pi^0\pi^0$  in the  $\pi^+\pi^-$  threshold region,  
 at the first order in an expansion in powers of  $a_i$

**The cusp singularity is proportional to  $a_0 - a_2$**

## Pions and QCD

The QCD Lagrangian

( mass terms)

( e.m. interactions)

$$\begin{aligned} \mathcal{L} = & (\bar{\psi}_L i\gamma \cdot \mathcal{D} \psi_L) + (\bar{\psi}_R i\gamma \cdot \mathcal{D} \psi_R) \\ & + (\bar{\psi}_R M \psi_L) + (\bar{\psi}_L M \psi_R) \\ & + e (\bar{\psi}_L Q \gamma \cdot A \psi_L) + e (\bar{\psi}_R Q \gamma \cdot A \psi_R) \end{aligned}$$

$M$  is the mass matrix and  $Q$  the charge matrix. Neglecting strange and heavier quarks,

$$M = \begin{vmatrix} m_u & 0 \\ 0 & m_d \end{vmatrix}, \quad Q = \begin{vmatrix} 2/3 & 0 \\ 0 & -1/3 \end{vmatrix}$$

In the limit  $M = 0, e = 0$  there is exact  $SU(2) \times SU(2)$  chiral symmetry — separate isospin for left handed and right handed quarks. In the real world this symmetry is broken to  $SU(2)$  — isospin, with pions acting as Nambu-Goldstone bosons. This leads to accurate predictions for the low energy pion interactions, both in  $\pi$ -Nucleon and in  $\pi - \pi$  scattering. In comparing with experimental data also take into account isospin breaking from  $m_d - m_u$  and radiative corrections.

These considerations can be extended to include the strange quark – and kaons. We will not discuss these extensions here.

## Pion scattering lengths and chiral dynamics

In the limit of exact I-spin (neglect electromagnetic interactions and the  $\pi^+ - \pi^0$  mass difference) we have accurate predictions for the  $\pi - \pi$  scattering lengths.

The scattering length defines the the low energy limit of the scattering amplitude.

$$\text{Weinberg 1966} \quad : \quad a_0 m_{\pi^+} = \frac{7 m_{\pi^+}^2}{16\pi f_\pi^2} = 0.159$$

$$a_2 m_{\pi^+} = \frac{-m_{\pi^+}^2}{8\pi f_\pi^2} = -0.045$$

$$\text{Colangelo et al. 2001} \quad : \quad a_0 m_{\pi^+} = 0.220 \pm 0.005$$

$$a_2 m_{\pi^+} = -0.0444 \pm 0.0010$$

$$(a_0 - a_2) m_{\pi^+} = 0.265 \pm 0.004$$

Can we match this precision?

## The problem of I-spin breaking — a pragmatic approach

The scattering length defines the scattering amplitude at threshold for S-wave states:

$$\mathcal{M}_{II} \approx \frac{8a_I m_\pi}{\pi} \quad (I = 0, 2)$$

We cannot measure directly  $a_0$  and  $a_2$ , which are defined in the unphysical limit of exact I-spin, but only the amplitudes for specific  $\pi\pi$  scattering channels at the respective thresholds. Identifying  $m_\pi$  with  $m_{\pi^+}$  we will thus *define* :

$\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$	$\mathcal{M}_{00} = \frac{8a_{00} m_{\pi^+}}{\pi}$	$(\pi^+ \pi^- \text{ threshold})$	$a_{00} \xrightarrow{\text{I-spin}} \frac{a_0 + 2a_2}{3}$
$\pi^+ \pi^0 \rightarrow \pi^+ \pi^0$	$\mathcal{M}_{+0} = \frac{8a_{+0} m_{\pi^+}}{\pi}$	$(\pi^+ \pi^0 \text{ threshold})$	$a_{+0} \xrightarrow{\text{I-spin}} \frac{a_2}{2}$
$\pi^+ \pi^- \rightarrow \pi^0 \pi^0$	$\mathcal{M}_x = \frac{8a_x m_{\pi^+}}{\pi}$	$(\pi^+ \pi^- \text{ threshold})$	$a_x \xrightarrow{\text{I-spin}} \frac{a_0 - a_2}{3}$
$\pi^+ \pi^- \rightarrow \pi^+ \pi^-$	$\mathcal{M}_{+-} = \frac{8a_{+-} m_{\pi^+}}{\pi}$	$(\pi^+ \pi^- \text{ threshold})$	$a_{+-} \xrightarrow{\text{I-spin}} \frac{2a_0 + a_2}{6}$
$\pi^+ \pi^+ \rightarrow \pi^+ \pi^+$	$\mathcal{M}_{++} = \frac{8a_{++} m_{\pi^+}}{\pi}$	$(\pi^+ \pi^+ \text{ threshold})$	$a_{++} \xrightarrow{\text{I-spin}} a_2$

## I-spin breaking in $\pi\pi$ threshold scattering

In order to compare with experimental measurements of  $\pi\pi$  scattering lengths chiral perturbation theory should include the effects of I-spin breaking. A first step (private communication by G. Colangelo, J. Gasser, B. Kubis and A. Rusetsky, to be considered preliminary) was the calculation of  $O(e^2)$  corrections at the lowest order in Chiral perturbation theory. The corrections are small but significant, e.g.


$$a_x = \frac{a_0 - a_2}{3} (1 + 2.2 \times 10^{-2})$$

so that the chiral perturbation theory predictions for  $a_x$  should become

$$a_x m_{\pi^+} = \frac{(0.271 \pm 0.004)}{3}$$

This is not the radiative correction, but only the chiral perturbation theory correction due to the electromagnetic  $m_{\pi^+} - m_{\pi^0}$  mass difference

# The Cusp in $K^+ \rightarrow \pi^+ \pi^0 \pi^0$



$\pi^+ \pi^- \rightarrow \pi^0 \pi^0$  re-scattering from  $K^+ \rightarrow \pi^+ \pi^+ \pi^-$  causes a cusp singularity in the  $\pi^0 \pi^0$  spectrum at the  $\pi^+ \pi^-$  threshold.

The cusp is proportional to  $(a_0 - a_2)$

N. Cabibbo, Phys. Rev. Lett. **93** (2004) 121801.

## How does the cusp arise

Let us write:

$$\mathcal{M}(K^+ \rightarrow \pi^+ \pi^0 \pi^0) = \mathcal{M} = \mathcal{M}_0 + \mathcal{M}_1$$

where  $\mathcal{M}_1$  is the contribution of the re-scattering graph.

We must consider two cases, with  $s_{\pi\pi}$  above or below the  $\pi^+ \pi^-$  threshold.

$$s_{\pi\pi} > 4m_{\pi^+}^2 : \quad \mathcal{M}_1 = i2a_x m_{\pi^+} \mathcal{M}_{+, \text{thr}} \sqrt{(s_{\pi\pi} - 4m_{\pi^+}^2) / s_{\pi\pi}}$$

$$|\mathcal{M}|^2 = (\mathcal{M}_0)^2 + |\mathcal{M}_1|^2$$


$$s_{\pi\pi} < 4m_{\pi^+}^2 : \quad \mathcal{M}_1 = -2a_x m_{\pi^+} \mathcal{M}_{+, \text{thr}} \sqrt{(4m_{\pi^+}^2 - s_{\pi\pi}) / s_{\pi\pi}}$$

$$|\mathcal{M}|^2 = (\mathcal{M}_0)^2 + (\mathcal{M}_1)^2 + 2\mathcal{M}_0\mathcal{M}_1$$

where  $\mathcal{M}_{+, \text{thr}}$  is the value of the  $K^+ \rightarrow \pi^+ \pi^+ \pi^-$  amplitude at the  $\pi^+ \pi^-$  threshold.

- Above threshold:  $\mathcal{M}_1$  imaginary — No interference
- Below threshold:  $\mathcal{M}_1$  real and negative — Interferes destructively with  $\mathcal{M}_0$

## How the Cusp looks



The  $\pi^0\pi^0$  spectrum in  $K^+ \rightarrow \pi^+\pi^0\pi^0$  in the  $\pi^+\pi^-$  threshold region,  
 at the first order in an expansion in powers of  $a_i$

## NA48 data and the cusp

With  $10^8 K^+ \rightarrow \pi^+ \pi^0 \pi^0$  events, a 1÷2 % measurement of  $(a_0 - a_2)$  seems possible, but since the cusp is a 10% effect, this **requires a theory good to 1 ÷ 2 parts in  $10^{-3}$ , and implicates higher order rescattering effects, and radiative corrections** .

It is possible to set up a systematic computation of the singular parts of an amplitude in terms of its non-singular parts. This leads to an expansion of the  $K \rightarrow 3\pi$  amplitudes in powers of the  $\pi\pi$  scattering lengths  $a_0, a_2$ .

$3\pi \rightarrow 3\pi$  scattering amplitudes are also implicated but give negligible correction at our target precision

The development is useful because the scattering lengths are small, which is a general consequence of the fact that pions act as “pseudo Goldstone Bosons” for chiral symmetry breaking.

The development to the second order in powers of  $a_0, a_2$  reveals **a second cusp** above the  $\pi^+ \pi^-$  threshold.

## New cusp results


With Gino Isidori we have completed a computation of the  $O(a_i^2)$  corrections to the  $K \rightarrow 3\pi$  amplitudes. This work includes:

- Other rescattering corrections:  $\pi^0\pi^0 \rightarrow \pi^0\pi^0$ ,  $\pi^+\pi^0 \rightarrow \pi^+\pi^0$ , etc.
- Corrections from “two-loops” graphs.
- It covers  $K^+ \rightarrow \pi^+\pi^0\pi^0$ ,  $K_L \rightarrow \pi^0\pi^0\pi^0$ , which have cusps, and  $K^+ \rightarrow \pi^+\pi^+\pi^-$ ,  $K_L \rightarrow \pi^+\pi^-\pi^0$ . which do not.

This work allows for a determination of  $a_0 - a_2$  with 5% theoretical uncertainty. To reach a 1% theoretical uncertainty we need:

- An evaluation of smaller (three loop) corrections.
- An evaluation of radiative corrections.

N. Cabibbo and G.Isidori, J. High Energy Phys. JHEP03(2005)021



$K \rightarrow 3\pi$  rescattering topologies at the two-loop level:

- a) single-channel  $\pi\pi$  scattering;
- b) irreducible  $3\pi \rightarrow 3\pi$  contributions;
- c)  $3\pi \rightarrow 3\pi$  amplitude due to  $\pi\pi$  scattering.

## Time reversal, Unitarity and Analyticity

The structure of the cusp is largely determined by very general principles.

1. Time reversal: In the sectors of interest the  $\mathbf{S}$ -matrix is symmetric,

$$\langle B|\mathbf{S}|A\rangle = \langle A|\mathbf{S}|B\rangle \quad K \rightarrow 3\pi \text{ and } \pi\pi \rightarrow \pi\pi$$

2. Unitarity:

$$\mathbf{S} = \mathbf{1} + i(\mathbf{R} + i\mathbf{I}) \quad \text{where } \mathbf{R} \text{ and } \mathbf{I} \text{ are hermitian.}$$

From time reversal:  $\mathbf{R}$  and  $\mathbf{I}$  are symmetric, their matrix elements are the real and imaginary parts of the matrix elements of  $\mathbf{S}$ .

$$2\mathbf{I} = \mathbf{R}^2 + \mathbf{I}^2 \quad \text{or, solving for } \mathbf{I},$$

$$\mathbf{I} = \mathbf{1} - \sqrt{\mathbf{1} - \mathbf{R}^2} = \frac{1}{2}\mathbf{R}^2 + \frac{1}{8}\mathbf{R}^4 + \frac{1}{16}\mathbf{R}^6 + \frac{5}{128}\mathbf{R}^8 \dots$$

3. Analyticity.

## $\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$ scattering at $\mathbf{O}(a_i^3)$ — 1

Define the “velocities”,

$$v_{\pm}(s) = \sqrt{\frac{|s - 4m_{\pi^+}^2|}{s}} \quad v_{00}(s) = \sqrt{\frac{|s - 4m_{\pi^0}^2|}{s}}$$

We can then write

$$\begin{aligned} \mathcal{M}_{00} &= A_{00} + B_{00} v_{\pm}(s) & s > 4m_{\pi^+}^2 \\ \mathcal{M}_{00} &= A_{00} + iB_{00} v_{\pm}(s) & s < 4m_{\pi^+}^2, \end{aligned}$$

and, for  $\pi^+ \pi^- \rightarrow \pi^0 \pi^0$ ,

$$\mathcal{M}_x = A_x + B_x v_{\pm}(s) \quad s > 4m_{\pi^+}^2$$

where  $A_{00}, B_{00}, A_x, B_x$  are regular (analytic) at the  $\pi^+ \pi^-$  threshold.

## $\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$ scattering at $\mathbf{O}(a_i^3)$ — 2

We can express  $\text{Re}(A)$  as a polynomial in  $s$ . Write:

$$\text{Re}(A_{00}) = \frac{8a_{00}(s)}{\pi}; \quad a_{00}(s) = a_{00} \left[ 1 + r_{00} \frac{(s - 4m_{\pi^+}^2)}{4m_{\pi^+}^2} + \dots \right]$$

and similarly for  $\pi^+ \pi^- \rightarrow \pi^0 \pi^0$ ,

$$\text{Re}(A_x) = \frac{8a_x(s)}{\pi}; \quad a_x(s) = a_x \left[ 1 + r_x \frac{(s - 4m_{\pi^+}^2)}{4m_{\pi^+}^2} + \dots \right]$$

In the limit of exact SU(2),  $a_{00} = \frac{a_0 + 2a_2}{3}$ ,  $a_x = \frac{a_0 - a_2}{3}$

The  $\pi^+ \pi^-$  intermediate state contributes to  $\text{Im} \mathcal{M}_{00}$  only above the  $\pi^+ \pi^-$  threshold, while the  $\pi^0 \pi^0$  state contributes both above and below, so that, at  $\mathbf{O}(\mathbf{R}^2)$ ,

$$\text{Im} \mathcal{M}_{00} = \frac{\pi}{4} v_{\pm}(s) (\text{Re} \mathcal{M}_x)^2 \Theta(s - 4m_{\pi^+}^2) + \frac{\pi}{8} v_{00}(s) (\text{Re} \mathcal{M}_{00})^2 + \mathbf{O}(\mathbf{R}^4)$$

## $\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$ scattering at $\mathbf{O}(a_i^3)$ — 3


Note that  $\text{Im } \mathcal{M}_{00} = \begin{cases} \text{Im } A_{00} + \text{Im } B_{00} \nu_{\pm}(s) & : \quad s > 4m_{\pi^+}^2 \\ \text{Im } A_{00} + \text{Re } B_{00} \nu_{\pm}(s) & : \quad s < 4m_{\pi^+}^2 \end{cases}$

Applying unitarity,  $\mathbf{I} = \mathbf{R}^2/2$  at  $\mathbf{O}(\mathbf{R}^2)$  both above and below the  $\pi^+ \pi^-$  threshold we then obtain

$$\begin{aligned} \text{Im } B_{00} &= \frac{\pi}{4} (\text{Re } A_x)^2 = \frac{16}{\pi} (a_x(s))^2 \\ \text{Re } B_{00} &= -\frac{\pi \nu_{00}(s)}{4} \text{Re } A_{00} \quad \text{Im } B_{00} = -\frac{32 \nu_{00}(s)}{\pi} a_{00}(s) (a_x(s))^2 \\ \text{Im } A_{00} &= \frac{\pi \nu_{00}(s)}{8} (\text{Re } A_{00})^2 = \frac{8 \nu_{00}(s)}{\pi} (a_{00}(s))^2 \end{aligned}$$


These results (and similar ones for other channels) are then fed into the unitarity relations for  $K \rightarrow 3\pi$  to obtain all corrections at  $\mathbf{O}(a_i^2)$ .

## The cusp at two loops



The theoretical  $\pi^0\pi^0$  spectrum in  $K^+ \rightarrow \pi^+\pi^0\pi^0$  (lower) computed at second order in powers of the scattering lengths. At the two-loops level also the “non-cusp” amplitude has an imaginary part arising from rescattering in  $\pi^+\pi^0 \rightarrow \pi^+\pi^0$ , etc.

## The cusp region



Theoretical  $\pi^0\pi^0$  spectrum in  $K^+ \rightarrow \pi^+\pi^0\pi^0$  (lower) at  $O(a_i^2)$   
normalized to the “unperturbed” amplitude.

The picture will be complicated by the existence of a  
 $\pi^+\pi^-$  ponium atom signal at the  $\pi^+\pi^-$  threshold!

## Space for improvement

- Stopping at  $O(a_i^2)$  level and neglecting radiative corrections implies a  $\sim 5\%$  theoretical error on  $(a_0 - a_2)$ . This may be sufficient at the present state of NA48 systematics.
- $O(a_i^3)$  terms and radiative corrections can be computed with a finite, but lengthy effort.
- Compute  $O(a_i^3)$  terms.
- Compute radiative corrections
- These computations will to a large extent be independent from Chiral Perturbation Theory which we wish to test!

# Conclusions

- The experimental study of  $K \rightarrow 3\pi$  decays is a powerful tool for gathering information on  $\pi\pi$  scattering in the low energy region.
- One can systematically evaluate rescattering effects in  $K \rightarrow 3\pi$  decays by means of an expansion in powers of the  $\pi\pi$  scattering lengths, and of other  $\pi\pi$  scattering parameters.
- This approach is less ambitious than the ordinary loop expansion performed in effective field theories, such as CHPT: the scope is not a dynamical calculation of the entire decay amplitudes, but a systematic evaluation of the **singular terms** due to rescattering effects.
- We have explicitly computed all the  $O(a_i^2)$  corrections to the leading cusp effect in  $K^+ \rightarrow \pi^+ \pi^0 \pi^0$ . The new terms produce a smaller square-root cusp behavior also above the  $\pi^+ \pi^-$  singularity.
- The 5% level of precision is probably not sufficient to fully exploit the potentially very accurate data of NA48, and is also larger than the error on the CHPT predictions of  $a_0 - a_2$ . To improve the situation we need a complete evaluation of the  $O(a_i^3)$  corrections and of the effects due to radiative corrections. Only first steps in this direction have been taken
- The theoretical error on  $(a_0 - a_2)$  can be reduced to the 1% level at both in theory and experiment. This might turn into the most accurate test of Chiral Perturbation Theory