


New results on J/ψ and ψ' nuclear absorption in p-A and S-U collisions at the CERN/SPS

Gonalo Borges, LIP-Lisbon


for the

NA50 Collaboration



- **Motivation: Why collect proton data?**
- **State of the art (QM2002)**
- **The experimental apparatus**
- **Analysis method**
- **J/ψ , ψ' and DY cross-sections at 400 GeV**
- **Comparison with previous p-A and light ion results**
- **Conclusions**

Motivation: Why collect proton data?



- NA50 studies J/ψ production in Pb-Pb collisions


In order to distinguish between

J/ψ absorption in nuclear matter & any J/ψ abnormal suppression

it is necessary to do a systematic study with several systems where it is assumed that only normal absorption plays a role.

- A good baseline is needed
↪ Such a baseline can be provided by proton-nucleus data.

The state of the art (QM2002)



- Data are plotted as a function of the average distance traveled by the $c\bar{c}$ pair through nuclear matter.

- A Simultaneous Glauber fit is performed using $\frac{\psi}{DY_{2.9-4.5}}$ results from

↪ p-p, p-d at 450 GeV

↪ p-A 98/00 at 450 GeV


↪ p-U, p-W at 200 GeV

↪ S-U at 200 GeV

- The line joins together the results of the best Glauber fit which leads to

$$\hookrightarrow \sigma_{\text{abs}}^{\psi} = 4.4 \pm 0.5 \text{ mb} \\ (\chi^2/dof = 1.0)$$

The experimental setup



J/ψ is detected via its decay into muon pairs $J/\psi \rightarrow \mu^+ \mu^-$

- Dimuon detection in:

$$2.92 < y_{\text{Lab}} < 3.92 \quad |\cos \theta_{\text{CS}}| < 0.5$$

- Proton beams at 450 or 400 GeV energy

↪ 3 independent detectors to measure the number of incident protons

- Several fixed targets: Be, Al, Cu, Ag, W, Pb

- Typical acceptances:

$$\hookrightarrow A^{J/\psi} = 13.8 \%, \quad A_{2.9-4.5}^{\text{DY}} = 14.5 \%, \quad A^{\psi'} = 16.3 \%$$

NA50 p-A runs

Data sample	Energy (GeV)	target thicknesses	targets	beam intensity (protons/s)	$N^{J/\psi}$ ($\times 10^3$)
1996-1998	450	26-39% λ_I	Be,Al,Cu,Ag,W	$(4 - 13) \times 10^8$	350 : 800
1998-2000	450	26-39% λ_I	Be,Al,Cu,Ag,W	$(0.8 - 2.5) \times 10^8$	80 : 180
2000	400	26-39% λ_I	Be,Al,Cu,Ag,W,Pb	$(9 - 13) \times 10^8$	38 : 68


- Absolute cross-sections from data collected in different years are affected by different systematic effects giving additional problems to the study of the **charmonia absorption using several target nuclei**
- To **minimize** such systematics, the last NA50 p-A run was taken
 - ↪ On a **very short period of time** at high intensity
 - ↪ **Using 6 targets** on a run by run **rotating regime**



The goal of the **last 2000 NA50 p-A run** was to precisely measure the **J/ψ absorption cross-section**

Analysis method

$$\frac{dN}{dM_{\mu\mu}} = N_{J/\psi} \frac{dN_{J/\psi}}{dM} + N_{\psi'} \frac{dN_{\psi'}}{dM} + N_{DY} \frac{dN_{DY}}{dM} + N_{D\bar{D}} \frac{dN_{D\bar{D}}}{dM} + \frac{dN_{Bkg}}{dM} + \frac{dN_{Empty}}{dM}$$



- **Opposite sign mass spectrum ingredients**

- ↪ $J/\psi \rightarrow \mu^+ \mu^-$

- ↪ $\psi' \rightarrow \mu^+ \mu^-$

- ↪ **Drell-Yan** ($q\bar{q} \rightarrow \mu^+ \mu^-$)

- ↪ **Simultaneous semi-leptonic decays of open charm mesons**

- ↪ **Combinatorial background**


$$N_{Bkg}^{+-} = 2R\sqrt{N^{++}N^{--}}$$

- **MC is treated as real data**

- ↪ **Acceptances and line shapes are determined via Monte Carlo and spectrometer simulation**

DY cross-sections at 400 GeV

$$\sigma_{p-A}^{\text{DY}} = \sigma_0^{\text{DY}} \times A^\alpha$$



- α^{DY} is expected to be 1 since DY production is **proportional** to the number of nucleon - nucleon collisions

- The fit to our data leads to

$$\alpha^{\text{DY}} = 0.986 \pm 0.018 \pm 0.008$$

($\chi^2/dof = 0.6$)

in nice agreement with the expectation.

- Nuclear effects (shadowing) are negligible in the explored phase space

J/ψ and ψ' absorption in nuclear matter



$$\sigma_{p-A}^{J/\psi, \psi'} = \sigma_0^{J/\psi, \psi'} \times A^\alpha$$

$$\alpha^{J/\psi} = 0.931 \pm 0.002 \pm 0.007$$

($\chi^2/dof = 1.4$)


$$\alpha^{\psi'} = 0.858 \pm 0.017 \pm 0.008$$

($\chi^2/dof = 2.2$)



★ Data show a **larger ψ' absorption** as compared to **J/ψ**

The Glauber model



↪ Charmonia production follows the hard process cross-section

$$\sigma_{p-A} = A \sigma_{NN}$$

↪ After production, charmonia states **can interact with the surrounding nuclear matter** with a given cross-section (σ_{abs}).

- Taking into account both processes

↪ production of the charmonium state,

↪ Possible absorption on it's way through nuclear matter, we get:


$$\frac{\sigma_{p-A}}{A} = \sigma_0 \frac{1}{(A-1)\sigma_{abs}} \times \int d^2b e^{-(A-1)T_A(\vec{b})} \sigma_{abs}$$

$T_A(\vec{b})$: Nuclear thickness function


- Charmonia experimental cross-sections can be fitted using this **Glauber model** with 2 free parameters:

$$\leftrightarrow \sigma_0, \sigma_{abs}$$

- J/ψ and ψ' results



$$\sigma_{\text{abs}}^{\psi} = 4.2 \pm 0.5 \text{ mb}$$
$$(\chi^2/dof = 0.9)$$



$$\sigma_{\text{abs}}^{\psi'} = 9.6 \pm 1.6 \text{ mb}$$
$$(\chi^2/dof = 2.0)$$

Comparison: All NA50 p-A data

- 3 NA50 p-A data samples:

↪ 96/98 data, High Intensity, 450 GeV [1]

↪ 98/00 data, Low Intensity, 450 GeV (average of analyses [1,2])

↪ 2000 data, 400 GeV

- Values from the different p-A samples

Individual $\sigma_{\text{abs, p-A}}$ Glauber fits

Data	\sqrt{s} (GeV)	$\sigma_{\text{abs}}^{\psi}$ (mb)	χ^2/dof	$\sigma_{\text{abs}}^{\psi'}$ (mb)	χ^2/dof
1996-1998 (HI)	29.1	4.8 ± 1.0	0.8	8.0 ± 1.4	1.2
1998-2000 (LI)	29.1	4.7 ± 1.0 (*)	1.0	6.0 ± 1.9	0.6
2000 (VHI)	27.4	4.2 ± 0.5	0.9	9.6 ± 1.6	2.0

- All results are **compatible** ↪ Perform **simultaneous fit**

(*) Estimated from $\frac{B_{\mu\mu}\sigma(\psi)}{\sigma(DY_{2.9-4.5})}$

[1] Euro Phys J C, in print.

[2] B. Alessandro et al. (NA50 Collaboration), Phys. Lett. B553 (2003) 167.

- **Method to estimate J/ψ absorption**

- ↳ Use the best estimate of each individual analysis ($\frac{\psi}{DY}$ or ψ)

- ↳ Fit all samples with a **common** σ_{abs}

- ↳ Leave normalizations as free parameters to take into account the different energies and kinematical conditions.

- **Method to estimate ψ' absorption**

- ↳ Use the best estimate of each individual analysis ($\frac{\psi'}{\psi}$)

- ↳ Fit all samples with a **common** σ_{abs}


- ↳ Assume that there is no $\frac{\psi'}{\psi}$ energy dependence.

- **For the last NA50 p-A data set**

- ↳ Luminosity systematic errors have been neglected since they affect all targets in essentially the same way (no effect on σ_{abs} measurement).

$\sigma_{\text{abs}}^{\psi}$

simultaneous fit in p-A data



- Perform a simultaneous Glauber fit

with $\frac{B_{\mu\mu}\sigma(\psi)}{\sigma(DY_{2.9-4.5})}$ results from :

↪ p-p, p-d at 450 GeV

↪ p-A 98/00 at 450 GeV

and $\frac{B_{\mu\mu}\sigma(\psi)}{A}$ results from:


↪ p-A 96/98 at 450 GeV

↪ p-A 2000 at 400 GeV

Glauber fit	$\sigma_{\text{abs}}^{\psi}$ (mb)	χ^2/dof
Without p-p, p-d	4.4 ± 0.4	0.8
With p-p, p-d	4.3 ± 0.3	0.7

A^{α} fit	α^{ψ}	χ^2/dof
Without p-p, p-d	0.929 ± 0.006	1.1
With p-p, p-d	0.941 ± 0.004	1.7

$\sigma_{abs}^{\psi'}$ simultaneous fit in p-A data



- Performing a simultaneous Glauber fit using $\frac{\psi'}{\psi}$ results from:

↪ p-A 98/00 at 450 GeV

↪ p-A 96/98 at 450 GeV

↪ p-A 2000 at 400 GeV

and assuming a J/ψ absorption

$$\sigma_{abs}^{\psi} = 4.4 \pm 0.4 \text{ mb}$$

we get

Glauber fit	$\sigma_{abs}^{\psi'}$ (mb)	χ^2/dof
	7.9 ± 0.6	1.0

$$\sigma_{abs}^{\psi'} - \sigma_{abs}^{\psi} = 3.5 \pm 0.7 \text{ mb}$$

$\sigma_{\text{abs}}^{\psi}$ from NA38 S-U published data

- NA38 has measured $\frac{B_{\mu\mu}\sigma(\psi)}{\sigma(DY_{2.9-4.5})}$ at 200 GeV as a function of E_T .

↪ The full Glauber model taking into account the centrality of the collision can be used to obtain the **J/ψ absorption in light ions collisions.**


↪ Published results show

$$\sigma_{\text{abs}}^{\psi}(S - U) = 7.1 \pm 3.0 \text{ mb}$$

- NA38 S-U data are now **reanalysed using the best of our present knowledge.**

↪ Same methods and procedures as used in Pb-Pb analyses

S-U reanalysis results



- **Reanalysis conditions:**
 - ↪ 6 different bins
 - ↪ MRS A (low Q^2) set of PDF
 - ↪ Same analysis procedure as used in NA50 PbPb data.
- **New absorption cross-section extracted from S-U collisions:**
$$\sigma_{abs}^{\psi} = 7.2 \pm 3.2 \text{ mb}$$
- **Error bar is due to the :**
 - ↪ Small Drell-Yan statistics
 - ↪ Large correlation between normalization and σ_{abs}^{ψ}


$$\sigma_{abs}^{\psi} \text{ (p-A)}$$

$$\sigma_{abs}^{\psi} \text{ (S-U)}$$

$$4.3 \pm 0.3 \text{ mb}$$

$$7.2 \pm 3.2 \text{ mb}$$

Simultaneous Glauber fit with p -A and S-U



- Perform a simultaneous Glauber fit

with $\frac{B_{\mu\mu}\sigma(\psi)}{\sigma(DY_{2.9-4.5})}$ results from :

- ↪ p - p , p - d at 450 GeV
- ↪ p -A 98/00 at 450 GeV
- ↪ S-U at 200 GeV

and $\frac{B_{\mu\mu}\sigma(\psi)}{A}$ results from:


- ↪ p -A 96/98 at 450 GeV
- ↪ p -A 2000 at 400 GeV

- New estimate of J/ψ absorption:

$$\sigma_{abs}^{\psi} = 4.3 \pm 0.3 \text{ mb}$$

$$(\chi^2/dof = 0.6)$$

Comparison with Pb-Pb results



- An absorption curve can be drawn as a function of E_T :
 - ↪ Using the Glauber model, the measured σ_{abs}^{ψ} and accounting for the different NA38 and NA50 calorimeter resolutions we can calculate the expected nuclear absorption for Pb-Pb collisions as a function of E_T .
 - ↪ The curve normalization is estimated from the S-U data rescaled to the Pb-Pb kinematical conditions.
 - ↪ The 2 curves represent the uncertainty on the absorption curve due to the $DY_{4.2-7.0}$ rescale from 200 to 158 GeV.

Conclusions

- Results from a new NA50 p-A data set, at 400 GeV incident energy, are now available
 - ↳ Systematics are minimized with rotating targets.
 - ↳ Data taking aimed for $\sigma_{\text{abs}}^{\psi}$ measurement.
- Global $\sigma_{\text{abs}}^{\psi}$ and $\sigma_{\text{abs}}^{\psi'}$ are obtained by means of a simultaneous fit using the best estimates of each NA50 p-A analyses

Global fit	$\sigma_{\text{abs}}^{\psi}$ (mb)	$\sigma_{\text{abs}}^{\psi'}$ (mb)	$\sigma_{\text{abs}}^{\psi'} - \sigma_{\text{abs}}^{\psi}$ (mb)
	4.3 ± 0.3	7.9 ± 0.6	3.5 ± 0.7

- Old NA38 S-U data were reanalysed and from it we obtain an absorption cross-section of $\sigma_{\text{abs}}^{\psi} = 7.2 \pm 3.2 \text{ mb}$.
- From a simultaneous fit including all NA50 p-A results and NA38 S-U reanalysis results we obtain $\sigma_{\text{abs}}^{\psi} = 4.3 \pm 0.3 \text{ mb}$.

Good precision in the J/ψ absorption cross-section measurement.

NA50 Collaboration Institutions

- **Università del Piemonte Orientale, Alessandria and INFN-Torino, Italy**
- **LAPP, CNRS-IN2P3, Annecy-le-Vieux, France**
- **LPC, Univ. Blaise Pascal and CNRS-IN2P3, Aubière, France**
- **IFA, Bucharest, Romania**
- **Università di Cagliari/INFN, Cagliari, Italy**
- **CERN, Geneva, Switzerland**
- **LIP, Lisbon, Portugal**
- **INR, Moscow, Russia**
- **IPN, Univ. de Paris-Sud and CNRS-IN2P3, Orsay, France**
- **Laboratoire Leprince-Ringuet, Ecole Polytechnique and CNRS-IN2P3, Palaiseau, France**
- **Università di Torino/INFN, Torino, Italy**
- **IPN, Univ. Claude Bernard Lyon-I and CNRS-IN2P3, Villeurbanne, France**
- **YerPhI, Yerevan, Armenia**