List of Tables

2.1 *Characteristics of the absorber and pre-absorber materials.* 47
2.2 *Targets characteristics for the data taking periods p-A 1996-1997.* . . 57

3.1 *Kinematic cuts imposed on the dimuon kinematic variables.* 63
3.2 E_T *binning used in the analysis of the PbPb and SU sets of data.* . . 68

4.1 *Values of the parameters of the Monte-Carlo DY fitting function.* . . 74
4.2 *Values of the parameters of the Monte-Carlo DY fitting function.* . . 74
4.3 *Total B.R. of charmed hadrons* H_c *in the semi-leptonic decay channel* ($\rightarrow \mu\overline{\nu}_{mu}X$)*[6].* 77
4.4 *Values of the parameters of the Monte-Carlo open charm fitting function.* 79
4.5 *Values of the parameters of the Monte-Carlo* J/ψ *fitting functions for the pA set-ups.* 84
4.6 *Values of the parameters of the Monte-Carlo* ψ' *fitting functions for the pA set-ups.* 84
4.7 *Values of the parameters of the Monte-Carlo* J/ψ *and* ψ' *fitting functions for the PbPb set-up.* 85
4.8 *Values of the coefficients* α *and* β *of the Monte-Carlo* J/ψ *and* ψ' *fitting functions.* 85
4.9 *Calculated acceptances (in %) for the indicated physical process.* . . . 88

5.1 *Characteristics of the materials crossed by pions and kaons when going through the hadron absorber.* 99
5.2 *Monte-Carlo* R_{BCK} *factor obtained for the 4 impact parameter b values chosen for the PbPb and SU systems.* 107
5.3 *Coefficients of the* 5^{th} *degree polynomials parametrising the b vs.* E_T *correlations for the PbPb and SU systems.* 110
5.4 *R factor for the pA systems. The high-intensity* R_{BCK} *values have been computed including the pile-up effect as explained in the previous paragraph. The pW 200GeV/c has been included to compare with the* R_{BCK} *factor obtained in the analysis of the two NA38 pW systems[3].* 113

5.5 R_{BCK} *factor for the 9* E_T *bins of the PbPb 158 A GeV/c system. The error on* R_{BCK} *comes from the fit to the 4 (+1 imposed for b=0 fm)* R_{BCK} *Monte-Carlo values vs. b.* 113
5.6 *R factor for the 5* E_T *bins of the SU 200 A GeV/c system. The error on* R_{BCK} *comes from the fit to the 4* R_{BCK} *Monte-Carlo values vs. b.* 113

6.1 $D\overline{D}/DY$ *obtained when fitting the 9 PbPb mass spectra with* $R_{BCK} = R_{BCK}^{MC}$. 123
6.2 $D\overline{D}/DY$ *obtained when the 9 PbPb opposite-sign mass distributions are fitted with the three different* $D\overline{D}$ *Monte-Carlo functionals corresponding to the 3 quoted values of* m_c. 124
6.3 $D\overline{D}/DY$ *obtained when fitting the SU opposite-sign mass spectra with the Monte-Carlo* R_{BCK} *values.* 124
6.4 $D\overline{D}/DY$ *values and* χ^2*'s obtained from the fit of the 9 PbPb mass spectra corresponding to the 9 PbPb* E_T *bins. The previous values are compared with the ones obtained using PYTHIA to generate all the Monte-Carlo shapes, with a Gaussian* k_T *distribution and* σ_{k_T} *= 0.9 GeV/c.* 138
6.5 *Isospin correction factors for the DY, calculated for 450 GeV/c pA(A=Al, Cu, Ag, W) interactions with the MRS A set of PDF and for M>1.5 GeV/*c^2*, -0.5<* $cos\theta$ *<0.5 and -0.52<* y *<0.48.* 139
6.6 $D\overline{D}/DY$ *obtained in the fit of the 9 PbPb mass spectra in the mass range 1.5<* M *<7.0 GeV/*c^2*, when the Monte-Carlo* R_{BCK} *values are used (0) and when* R_{BCK}*=1.0 for all the* E_T *bins (1).* 145
6.7 $D\overline{D}/DY$ *obtained in the fits of the SU opposite-sign mass spectra when the Monte-Carlo* R_{BCK} *are used (0) and when* R_{BCK}*=1.000 for all the* E_T *bins (1).* 147
6.8 *The Monte-Carlo nominal* R_{BCK} *values (0) are compared with the ones obtained when simultaneously fitting the 4 pA systems in the mass range 1.3<* M *<2.0 and 4.5<* M *<8.0 GeV/*c^2 *whilst leaving the* R_{BCK} *factors as free parameters in the fit (f).* 149

7.1 *Cross sections calculated for the three studied systems at the related energies. The four different cross section values listed for each of the three systems correspond to : B.R.(*$D \rightarrow \mu$*) = 1 and no.kin.cuts (p-p), true B.R.(*$D \rightarrow \mu$*) and no.kin.cuts (p-p B.R.) true B.R.(*$D \rightarrow \mu$*) and NA50 k.c (p-p B.R., k.c.) true B.R.(*$D \rightarrow \mu$*) and NA50 k.c and nuclear effects (A-B B.R., k.c.).* 158
7.2 *Quantities needed to calculate the luminosity for the PbPb, SU and pW systems : measured* J/ψ *cross section* $\sigma_{J/\psi}^{meas.}$*, number of* J/ψ *measured in the fit to the indicated systems and* J/ψ *acceptance. The calculated luminosities are also listed.* 159

7.3 *Quantities needed to compute the expected number of dimuons from open charm decay for the PbPb, SU and pW systems and for* m_c*=1.5 GeV/*c^2*. The listed number of dimuons from DY is necessary in order to calculate the ratio* $D\overline{D}/DY$ *which is also given.* 160
7.4 *Enhancement of the open-charm dimuon yield with respect to pA results for the pW, PbPb and SU systems. The enhancement E has been calculated for each of the considered* E_T *bin; the number of participant nucleons in the collision is also listed. The errors on E come from the error on the measured* $D\overline{D}/DY$ *(*$\simeq$*10%) and from the error on the calculation of the luminosity (7÷10%). The systematic errors due to the normalisation to the* $E\,|_{pA}$ *value, 8%, and to the 12% incertitude in the B.R. measurement, have not been included because they would affect all the listed values in the same way (and direction).* 163