

ISSN 1810-5408

# Nuclear Science and Technology

**Volume 9, Number 2, June 2019**

Published by

VIETNAM ATOMIC ENERGY SOCIETY

VIETNAM ATOMIC ENERGY INSTITUTE

# NUCLEAR SCIENCE AND TECHNOLOGY

Volume 9, Number 2, June 2019

## Editorial Board

### Editor-in-chief

Tran Huu Phat (VINATOM)

### Executive Editors

Vuong Huu Tan (VARANS)

Tran Chi Thanh (VINATOM)

Cao Đình Thanh (VINATOM)

Hoang Anh Tuan (VAEA)

### Editors

Nguyen Kien Cuong (VINATOM)

Nguyen Nhi Dien (VINATOM)

Nguyen Thi Kim Dung (VINATOM)

Ho Manh Dung (VINATOM)

Nguyen Nam Giang (VINATOM)

Trinh Van Giap (VINATOM)

Le Ngoc Ha (108 Military Central Hospital)

Phan Son Hai (VINATOM)

Le Huy Ham (VAAS)

Nguyen Quoc Hien (VINATOM)

Le Van Hong (VINATOM)

Nguyen Tuan Khai (VINATOM)

Pham Dinh Khang (VINATOM)

Le Hong Khiem (IOP)

Dao Tien Khoa (VINATOM)

Tran Hoai Nam (Duy Tan University)

Dang Duc Nhan (VINATOM)

Nguyen Hao Quang (VINATOM)

Nguyen Mong Sinh (VINATOM)

Tran Duc Thiep (IOP)

Dang Quang Thieu (VINATOM)

Le Ba Thuan (VINATOM)

Nguyen Trung Tinh (VARANS)

Tran Ngoc Toan (VINATOM)

Duong Thanh Tung (VARANS)

Nguyen Nu Hoai Vi (VARANS)

### Science Secretary

Hoang Sy Than (VINATOM)

---

**Copyright:** ©2008 by the Vietnam Atomic Energy Society (VAES), Vietnam Atomic Energy Institute (VINATOM).

### Published by

Vietnam Atomic Energy Society, 59 Ly Thuong Kiet, Hanoi, Vietnam

Tel: 84-24-39420463 Fax: 84-24-39424133 Email: nuscitech@vinatom.gov.vn

Vietnam Atomic Energy Institute, 59 Ly Thuong Kiet, Hanoi, Vietnam

Tel: 84-24-39420463 Fax: 84-24-39422625 Email: infor.vinatom@hn.vnn.vn

---

## Contents

---

On Burnup Modelling Issues Associated with VVER-440 Fuels <i>Branislav Vrban, Štefan Čerba, Jakub Lüley, Filip Osuský, Mikuláš Vorobel, Vladimír Nečas</i> .....	01
Estimation of Tritium Production in VVER-440 Reactor Core During Normal Operation <i>Štefan Čerba, Jakub Lüley, Branislav Vrban, Filip Osuský, Vladimír Nečas</i> .....	10
Processing of the multigroup cross-sections for MCNP calculations <i>Jakub Lüley, Branislav Vrban, Štefan Čerba, Filip Osuský, Vladimír Nečas</i> .....	17
Conceptual design of a small-pressurized water reactor using the AP1000 fuel assembly design <i>Van Khanh Hoang, Viet Phu Tran, Van Thin Dinh, Hoai Nam Tran</i> .....	25
Low-energy experiments at the S3 spectrometer <i>S. Franchoo</i> .....	31
Dosimetric characteristics of 6 MV photons from TrueBeam STx medical linear accelerator: simulation and experimental data <i>N. D. Ton, B. D. Linh, Q.T. Pham</i> .....	37
Conceptual designing of a slow positron beam system using Simion simulation program <i>Cao Thanh Long, Huynh Dong Phuong, Nguyen Trung Hieu, Tran Quoc Dung</i> .....	45

# Low-energy experiments at the S3 spectrometer

S. Franchoo

*IPN, 91406 Orsay, France*

*E-mail: franchoo@ipno.in2p3.fr*

(Received 08 November 2019, accepted 19 November 2019)

**Abstract:** With the advent of the Spiral-2 linear accelerator and the associated S3 spectrometer at the Ganil laboratory in France, new realms of intermediate-mass  $N=Z$  and very heavy nuclei will soon become available for research in nuclear structure. After their production and selection in the spectrometer, the ions of interest will be stopped in a buffer gas, neutralised, resonantly re-ionised, cooled and bunched. This will bring us in an adequate position to perform laser spectroscopy, mass measurements and decay spectroscopy of their ground and isomeric states. In this contribution we report on the ongoing commissioning of the detector set-up.

**Keywords:** *New facilities, laser spectroscopy, mass spectrometry, nuclear structure*

## I. INTRODUCTION

Recent advances in nuclear theory are bringing ever heavier nuclei within reach of ab-initio techniques. The quantum many-body calculations naturally include three-nucleon forces, emphasising their influence for instance in the Gamow-Teller strength of  $^{100}\text{Sn}$  [1]. Large-scale shell-model calculations are carried out in valence spaces that span complete shells and beyond, taking into account correlations that previously had to be left out. These in turn induce quadrupole deformation throughout the nuclear chart, affecting even the immediate vicinities of doubly magic nuclei such as  $^{78}\text{Ni}$  [2]. The Monte-Carlo shell model puts in place a framework for exploring the sudden quantum phase transitions that are observed in exotic isotopes. Striving to unite single-particle and collective behaviours, it stresses the role of the central and tensor forces in the shell evolution of magic and near-magic nuclei such as zirconium, tin and mercury [3]. Reaction theory increasingly embraces structure models. By replacing both the optical and the shell-model potential by the nucleon self-energy, the dispersive optical model links energy states

above and below the Fermi energy in a single description [4].

Experimentally, proton-induced fission at Isolde and Triumf gives access to the neutron-rich side of the nuclear chart. The particles diffuse out of the target as neutral atoms, they are ionised, extracted at low energy and eventually post-accelerated. At NSCL and Riken, fragmentation at high energy covers a broad spectrum of nuclei that are available for multinucleon transfer and knock-out reactions. In-flight spectroscopy of the emitted radiation relies on the development of multidetector arrays that provide the necessary granularity for event reconstruction. Fusion-evaporation reactions favour products rich in protons, recoiling into a spectrometer at threshold energies but setting aside for FLNR and S3 a range of experiments that would be difficult to perform elsewhere.

The S3 spectrometer for exotic radioactive ion beams is currently being installed at the Ganil laboratory in France and shall become operational in 2023. It offers novel opportunities to investigate isotopes produced in heavy-ion fusion-evaporation reactions, in particular for channels with low

cross sections at the proton dripline and near the upper end of the nuclear chart. An approach based on laser spectroscopy in a swiftly expanding gas jet is adopted for the detection strategy, called Radioactive Elements in a Gas jet for Laser Ionisation and Spectroscopy (Reglis). It shall be coupled to a mass spectrometer that is built around a linear ion trap, referred to as Piège à Ions Linéaire du Ganil pour la Résolution des Isobares et la mesure de Masse (Pilgrim).

## II. REPORT

### A. Physics scope

The Reglis experiment focusses on two objectives, the isotopes near the  $N=Z$  line from  $Z=40$  onwards and the very heavy nuclei with  $Z \geq 89$ . The first region allows to cover a broad range of physics issues that spans from collectivity and deformation over spin-aligned coupling schemes in nuclei such as  $^{90}\text{Rh}$  and  $^{94}\text{Ag}$ . Also the balance between isoscalar and isovector pairing interactions in  $^{98}\text{In}$  and the evolution of single-particle structures in  $^{99}\text{In}$  and  $^{101}\text{Sn}$  attracts much interest. We shall shed light on quadrupole moments in the tin chain and investigate novel decay modes such as super-allowed  $\alpha$ -decay in  $^{112}\text{Ba}$ - $^{108}\text{Xe}$ - $^{104}\text{Te}$  and cluster radioactivity in  $^{114,112}\text{Ba}$ . The measurement of masses will give access to binding energies and the role of the Wigner term. Among the heavy elements, we shall focus on the actinides. This includes the development of the  $N=126$  shell closure in  $^{210-213}\text{Ac}$  and  $^{213-215}\text{Th}$ , the appearance of octupole deformation in  $^{225-228}\text{U}$ , and the possible emergence of a superheavy spherical shell gap as seen through K-isomers in  $^{249-252}\text{Fm}$  and  $^{252-254}\text{No}$ .

The mean-square charge radii, magnetic dipole and electric quadrupole moments can be extracted from the hyperfine interaction

between the nucleus and its surrounding electrons, which is recorded by laser spectroscopy. The masses, spins and decay properties of their ground states and isomeric levels will be constrained by complementary measurements on the collected ions by means of time-of-flight mass spectrometry and decay spectroscopy.


### B. The S3 spectrometer

A new superconducting linear accelerator has been constructed at Ganil, comprising 26 accelerating cavities housed in 19 cryomodules [5]. It will be able to deliver stable ion beams with intensities that are ten times larger than what is presently available. With a mass-to-charge ratio of 3, currents of 3  $\mu\text{A}$  for  $^{40}\text{Ar}$ , 2  $\mu\text{A}$  for  $^{40}\text{Ca}$  and 1  $\mu\text{A}$  for  $^{48}\text{Ca}$  and  $^{58}\text{Ni}$  at energies up to 14.5 MeV per nucleon are projected. With a later upgrade of the injector to a mass-to-charge ratio of 7, heavy beams up to uranium will become available.

These primary beams will make it possible to study fusion-evaporation reactions with cross sections down to picobarns, yielding neutron-deficient nuclei near the  $N=Z$  line and isotopes of very heavy elements. They will be separated from the intense background in the S3 recoil spectrometer [6]. The spectrometer consists of two stages, the first one of which is a momentum achromat that lets the reaction products pass through while the primary beam is suppressed with a factor of  $10^3$ . Subsequently the ions of interest are selected in a mass separator with a resolution of 300. The combination of the momentum achromat and the mass separator results in a final rejection power of  $10^{13}$ . An angular acceptance of  $\pm 50$  mrad, a magnetic rigidity acceptance of  $\pm 7$  % and a charge-state acceptance of  $\pm 10$  % are aimed at. This means two charge states on either side of the centrally chosen charge state

will be transmitted. A high-resolution mode that separates the charge states as well as a converging mode with a better transport efficiency at the price of a wider beam spot are proposed. To illustrate with an example, for the  $^{208}\text{Pb}(^{48}\text{Ca},2n)^{254}\text{No}$  reaction at 217 MeV a transmission efficiency of 56% is estimated for 5 charge states within a circular area of 50 mm

diameter. The  $^{254}\text{No}$  isotopes arrive with a mean energy of 37 MeV. In this mass region the ion rate amounts to 1 particle per second only but the beam would be almost 100% pure. For the  $^{116}\text{Sn}(^{40}\text{Ar},4n)^{152}\text{Er}$  reaction at 167 MeV, on the other hand, the expected rate reaches  $8.6 \cdot 10^4$  particles per second yet with a purity of 1% only.



**Fig. 1.** Drawing of the Reglis gas cell. The S3 beam arrives from the left and the gas enters from below (orange). The stopped particles (red) are evacuated on the top right side

### C. Principle of the detector

To take the best advantage of the capabilities of the accelerator and the spectrometer, a detection set-up with high efficiency is in order. The ions should be made available for several measurement set-ups so they should be transferable as a beam at low emittance. Moreover and in spite of the  $10^{13}$  rejection power of S3, some secondary beams will still be significantly contaminated by their  $A \pm 1$  neighbours and additional selectivity is desirable.

We therefore aim to stop the reaction products in a buffer gas at high pressure [7]. They will be neutralised through collisions with the gas atoms, extracted from the gas cell in a well-defined cold and supersonic jet and resonantly re-ionised by laser light. They will then be sent through a series of radiofrequency

quadrupoles (RFQ), in which their emittance is greatly reduced before being fed into various experimental set-ups. At the same time, we shall be able to instantaneously perform laser spectroscopy in the gas jet by scanning the resonance wavelength across the hyperfine structure. This is possible because the low density in the uniform jet keeps pressure broadening small such that sufficient resolution can be achieved.

### D. Intrajet laser spectroscopy

The thickness of the entrance window is of the utmost importance as some of the secondary beams carry energies below as little as 1 MeV per nucleon. For critical cases we can afford not more than a couple of micrometers of material. On the other hand the window should span a diameter of 50 mm and remain rigid enough to withstand the pressure

difference with the high vacuum inside S3. We choose a titanium foil that is supported by a honeycomb grid, both fused together by friction stir welding.

For the buffer gas a compromise is sought between the stopping power in the gas cell and the load on the pumping system. We use argon at 200 to 500 mbar purified to the part-per-billion level. Special care is taken for the mechanical design of the cell so we obtain a gas flow that is as laminar as possible, avoiding turbulences that would trap the particles. The flow transports the radioactive particles to the exit hole with an evacuation time of several hundreds of milliseconds, setting a limit for the shortest lifetimes we can efficiently measure. The exit hole takes the shape of a Laval nozzle. The thermodynamic properties of the jet are defined by the background pressure in the experimental chamber as well as the contour of the nozzle, the machining of which demands a dedicated effort with a precision of 5  $\mu\text{m}$ . Several nozzles were tested, the latest design producing a homogeneous supersonic jet of Mach 8 at a temperature of 18 K over a distance of 25 mm. For development purposes the jet was visualised with Planar Laser-Induced Fluorescence and modelled with the Comsol and Ansys flow codes [8].

Laser ionisation happens in two or more steps. The last step may not be resonant and simply allow one of the atomic electrons to reach the continuum. One of the resonant steps is used for probing the hyperfine structure and requires a narrow bandwidth. It irradiates the gas jet perpendicularly and is expanded into a laser sheet to cover as large a section of the jet as possible. It is planned to implement a system of both solid-state as well as dye lasers at high repetition rate, the former being relatively free of safety constraints and the

latter providing higher power levels. From a proof-of-principle experiment that was performed on  $^{214-215}\text{At}$  at an overall efficiency of 10% and a selectivity of 3000 appears within reach [9]. The FWHM resolution of the hyperfine spectrum came down to 400 MHz during this measurement, but at S3 we intend to achieve 100 MHz, which will be essentially due to the gas temperature, the jet divergence and the laser linewidth. This number can be compared to the natural linewidth of the atomic transition of 4 MHz but remains amply sufficient to probe the hyperfine structures, the splittings of which typically amount to 300 MHz in the actinides.


### **E. Beam manipulation and mass spectrometry**

The ions that emerge from the supersonic jet will be captured in a S-RFQ. The name derives from its geometrical shape, the purpose of which is to move the ions vertically out of the axis along which the broadband laser is sent in. A quadrupole mass filter (QMF) with a resolving power of 100 further cleans the beam. It is followed by a RFQ-buncher, which is filled with  $10^{-2}$  to  $10^{-3}$  mbar of helium and cools and prepares the beam for injection into a drift tube. The tube lifts the ions to 3 kV and transfers them to the Pilgrim multireflection time-of-flight mass spectrometer (MR-tof-MS) [10]. The latter basically comprises a linear trap with an electrostatic mirror on either end, between which the particles travel back and forth until they are spatially separated from isobaric contaminants because of their mass differences. A resolution of  $10^5$  can be achieved for an input emittance of 10  $\pi\text{.mm.mrad}$ , corresponding to a flight path of some hundreds of meters in 10 ms. For a half-life of 100 ms, Pilgrim is expected to attain a sensitivity of one particle per ten seconds.

While the transmission through S3 can be evaluated at 50% on average, the thermalisation, neutralisation and transport of the particles in the gas cell are estimated at more than 25%. The rate of laser ionisation depends on the availability of a suitable scheme and can be taken at 50%, the capture efficiency in the RFQ sequence at 80%. The overall efficiency of our set-up before injection into Pilgrim would then reach at least 5%.

Reglis and Pilgrim are presently installed at the LPC laboratory nearby Ganil in Caen, where they can be commissioned over

the next two years. Stable beams will be produced by heating a metallic filament that is mounted inside the gas cell. In order to minimise adverse effects on the gas flow, the heat excess is removed by water cooling. One of the first elements to extract will be erbium, which is the chemical homologue of fermium. Since laser-ionisation schemes are poorly known for the actinides we are interested in at S3, they will be developed from better studied transitions in the lanthanides. According to the current timeline, S3 should become available for experiments in 2023.



**Fig. 2.** The Reglis and Pilgrim set-ups as they are presently installed at the LPC laboratory in Caen for commissioning

### F. Decay spectroscopy

The ions that emerge from Pilgrim will be implanted in a thin carbon foil that is surrounded by a small but segmented silicon array. Simple ion counting as well as decay spectroscopy through electron and  $\alpha$ -radiation will be handled by a new generation of

electronics with automatic gain switching. Particular attention will be paid to combining a high efficiency and resolution with a low energy threshold. Two germanium detectors can be integrated in a compact geometry for registering coincident  $\gamma$ -rays. The addition of a tape station for  $\beta$ -decay studies, which is



helpful for those physics cases at  $N=Z$ , is also foreseen.

### III. CONCLUSIONS

The new S3 spectrometer at Ganil shall unlock new  $N=Z$  nuclei and very heavy isotopes for experimental research. Because of the extraordinary low cross sections to produce these nuclei, a dedicated detector set-up that is sensitive to count rates of a few particles per second has been constructed and is now being commissioned. The Reglis gas cell provides the base for laser spectroscopy in a supersonic jet, probing the hyperfine structure to yield charge radii and electromagnetic moments. The Pilgrim trap is properly suited for mass measurements, recording the time of flight as the ion path is reflected between two electrostatic mirrors. In order to overcome the slow evacuation time of the gas cell, which constitutes the main limitation of our experiment when accessing the shortest lived isotopes, the development of a smaller gas volume that incorporates electrical radiofrequency fields is being considered.

### ACKNOWLEDGMENTS

The Reglis and Pilgrim experiments have been conceived and built and are since operated by a collaboration of the Ganil, IPN-Orsay, Irfu-Saclay, and LPC-Caen laboratories and the universities of Jyväskylä, Leuven, and Mainz. We are grateful to the many administrators, technicians, engineers, and physicists who have made the project possible. We thank the Programme blanc of the Agence national de recherche for its essential part in the funding of Reglis and the Brix network within the Interuniversity Attraction Poles of the Belgian Science Policy Office for its contribution to Pilgrim. We are indebted to the

Institut national de physique nucléaire et de physique des particules for continuous support.

### REFERENCES

- [1]. P. Gysbers et al., Discrepancy between experimental and theoretical  $\beta$ -decay rates resolved from first principles. *Nature Physics*, in print, 2019.
- [2]. F. Nowacki et al., Shape Coexistence in  $^{78}\text{Ni}$  as the Portal to the Fifth Island of Inversion. *Physical Review Letters* **117**, 272501 (2016).
- [3]. T. Togashi et al., Quantum Phase Transition in the Shape of Zr isotopes. *Physical Review Letters* **117**, 172502, 2016.
- [4]. M. Mahzoon et al., Forging the Link between Nuclear Reactions and Nuclear Structure. *Physical Review Letters* **112**, 162503 (2014).
- [5]. Ghribi et al., Status of the Spiral-2 linac cryogenic system. *Cryogenics* **85**, 44, 2017.
- [6]. F. Déchery et al., The Super Separator Spectrometer S3 and the associated detection systems. *Nuclear Instruments and Methods B* **376**, 125, 2016.
- [7]. R. Ferrer et al., In-gas laser ionization and spectroscopy experiments at the Superconducting Separator Spectrometer: Conceptual studies and preliminary design. *Nuclear Instruments and Methods in Physics Research B* **317**, 570, 2013.
- [8]. Zadornaya et al., Characterisation of Supersonic Gas Jets for High-Resolution Laser-Ionisation Spectroscopy of Heavy Elements. *Physical Review X* **8**, 041008, 2018.
- [9]. R. Ferrer et al., Towards high-resolution laser ionization spectroscopy of the heaviest elements in supersonic gas jet expansion. *Nature Communications* **8**, 14520, 2017.
- [10]. P. Chauveau et al., Pilgrim, a Multi-Reflection Time-of-Flight Mass Spectrometer for Spiral2-S3 at Ganil. *Nuclear Instruments and Methods in Physics Research B* **376**, 211, 2016.

---

# INSTRUCTIONS FOR AUTHORS

---

## GENERAL INFORMATION

Nuclear Science and Technology (NST), an international journal of the Vietnam Atomic Energy Society (VAES) and Vietnam Atomic Energy Institute (VINATOM), quarterly publishes articles related to theory and application of nuclear science and technology. All papers and technical notes will be refereed.

It is understood that the paper has been neither published nor currently submitted for publication elsewhere. The copyright of all published papers and notes will be transferred in VAES.

## DETAILED FIELDS

NST covers all fields of nuclear science and technology for peaceful utilization of nuclear energy and radiation. Authors should choose one of the following fields at the time they submit their manuscript: 1) Nuclear Physics, 2) Nuclear Data, 3) Reactor Physics, 4) Thermal Hydraulics, 5) Nuclear Safety, 6) Nuclear I&C, 7) Nuclear Fuel and Materials, 8) Radioactive Waste Management, 9) Radiation Protection, 10) Radiation Technology, 11) Nuclear Techniques in Food and Agriculture, 12) Nuclear Medicine and Radiotherapy, 13) Nuclear Techniques in Industries, 14) Environment Radioactivity, 15) Isotope Hydrology, 16) Nuclear Analytical Methods, 17) Health Physics, 18) Fusion and Laser Technology.

## MANUSCRIPT SUBMISSION

Manuscript for publication should be submitted to the Editorial Office in triplicate by postal mail. For electronical submission use [nuscitech@vinatom.gov.vn](mailto:nuscitech@vinatom.gov.vn).

### Submission Address

Department of Planning, R&D Management  
Vietnam Atomic Energy Institute, 59 Ly Thuong  
Kiet Street, Hanoi, Vietnam  
E-mail: [nuscitech@vinatom.gov.vn](mailto:nuscitech@vinatom.gov.vn).

## MANUSCRIPT PREPARATION

Manuscripts must be written in English with adequate margins and indented paragraph. All manuscript must use SI (metric) units in text, figures, and tables. Manuscripts should in general be organized in the following order: title, names of authors and their complete affiliation including zip code, abstract (not exceeding 200 words), keywords (up to 7), introduction, main body of a paper, acknowledgments, references, appendices, table & figure captions, tables and figures. Unnecessary sections may be omitted.

**Headings:** Use I, II, ... for major headings and A, B, ... for secondary headings.

**Mathematical formulas:** All mathematical formulas should be clearly written, with special consideration to distinctive legibility of sub- and superscripts. Equation (at least the principal ones) should be numbered consecutively using Arabic numerals in parentheses in the right hand margin.

**Tables and Figures:** Tables should be numbered with Roman numerals. Figures should be numbered consecutively with Arabic numerals in order of their first appearance and have a complete descriptive title. They should be typed on separate sheets. Tables should not repeat data which are available elsewhere in the paper. Figures should be original ink drawing or computer drawn figures in the original and of high quality, ready for direct reproduction. Figures should be referred to in the text as, for example, Fig. 1., or Fig. 2. .

**Reference:** References should be listed at the end of the text and presented as follows:

- [1] C. Y. Fu et al., *Nuclear Data for Science and Technology*, S. M. Qaim (Ed.), p. 587 (1991).
- [2] C. Kalbach, *Z. Phys.*, A283, 401 (1977).
- [3] S. Shibata, M. Imamura, T. Miyachi and M. Mutou, "Photonuclear spallation reactions in Cu", *Phys. Rev. C* 35, 254 (1987).

# KHOA HỌC VÀ CÔNG NGHỆ HẠT NHÂN

---

**Chịu trách nhiệm xuất bản**

TRẦN HỮU PHÁT

**Chịu trách nhiệm nội dung**

TRẦN HỮU PHÁT

TRẦN CHÍ THÀNH

**Trình bày**

LÊ THÚY MAI

---

In 200 cuốn, khổ 19x26,5cm tại Công ty TNHH Trần Công  
Địa chỉ: số 12 ngách 155/176 Đường Trường Chinh, Hà Nội  
Giấy đăng ký kế hoạch xuất bản số: 770/GP-BTTTT cấp ngày 20 tháng 5 năm 2011  
In xong và nộp lưu chiểu Quý IV năm 2019