


Standard Model Results from LEP

Pat Ward

University of Cambridge

- Introduction
- Electroweak Physics
- QCD and Two-photon Results



Introduction

LEP1	1989–1995	$\sqrt{s} \sim m_Z$	4.5M Z events / expt
LEP2	1996–2000	$161 < \sqrt{s} < 209 \text{ GeV}$	10k WW events/expt

- LEP was shut down at the end of 2000, so why are we still giving LEP talks?
- Many physics results still being produced
- In the last 2 years, the 4 LEP experiments (ALEPH, DELPHI, L3 and OPAL) have submitted for publication >100 papers
About 25% of these were on LEP1 physics
- The 4 expts have submitted > 120 papers to ICHEP04
Majority on Standard Model physics
Mostly final results, but some new preliminary results too
- This talk will concentrate on results which have been finalized, or are new, in the last year (but include some older important results too)

Introduction

LEP Standard Model physics covers a wide range of topics; no time in this talk for, e.g.,

- Precise measurement of the τ lifetime (DELPHI)
 $\tau_\tau = 290.9 \pm 1.4 \pm 1.0 \text{ fs}$ (c.f. PDG2004: $290.6 \pm 1.1 \text{ fs}$)
- τ branching ratios and strange spectral functions (DELPHI, OPAL)
- $\mu^+ \mu^-$, $\tau^+ \tau^-$ and hard-photon production in two-photon collisions (DELPHI, L3, OPAL)
- Excited b-hadrons, $B_s^0 - \overline{B}_s^0$ oscillations, B spectral moments (DELPHI)

Will give a brief overview of current results; for more details consult the experiments' web pages: <http://aleph.web.cern.ch/aleph/>

<http://delphiwww.cern.ch/>

<http://l3.web.cern.ch/l3/>

<http://opal.web.cern.ch/Opal/PPwelcome.html>

LEP1 Electroweak Physics

- Z lineshape measurements final since summer 2000

$$m_W = 91.1875 \pm 0.0021 \text{ GeV}$$


$$\Gamma_Z = 2.4952 \pm 0.0023 \text{ GeV}$$

$$\sigma_h^0 = 41.540 \pm 0.037 \text{ nb}$$

$$R_\ell = 20.767 \pm 0.025$$

$$A_{\text{FB}}^{0,\ell} = 0.01714 \pm 0.00095$$

- New measurement of A_{FB}^b (DELPHI)
- \Rightarrow new LEP heavy-flavour combination



$$A_{\text{FB}}^{0,b} = 0.0998 \pm 0.0017$$

LEP1 Electroweak Physics

- **OPAL** have published new measurements of Γ_u, Γ_d
- Use hadronic events with FSR ($Z \rightarrow q\bar{q}\gamma$)

Enriched in up-type quarks

$$\Gamma_{\text{had}} = 2\Gamma_u + 3\Gamma_d$$


$$\Gamma_{\text{had}+\gamma} \sim 8\Gamma_u + 3\Gamma_d$$

- Results:

$$\Gamma_u = 300_{-18}^{+19} \text{MeV}$$


$$\Gamma_d = 381_{-12}^{+12} \text{MeV}$$

- Good agreement with SM



- More precise than earlier measurements (**DELPHI, L3, OPAL**)


Two-fermion Production at LEP2



- Measure cross-sections and asymmetries for inclusive and 'non-radiative' events
- LEP combination of preliminary measurements

Good agreement with SM

⇒ limits on new physics, e.g. Z' , leptoquarks, RPV squarks, contact interactions, extra dimensions



Two-fermion Production at LEP2


- At LEP2 energies, γ -exchange becomes important \Rightarrow can measure γ -Z interference \Rightarrow almost model-independent determination of m_Z in S-matrix fit

- e.g. DELPHI results


$$m_Z = 91.1831 \pm 0.0034 \text{ GeV}$$

c.f. $m_Z = 91.1863 \pm 0.0028 \text{ GeV}$
from standard LEP1 fit assuming SM interference

- OPAL, DELPHI have finalized two-fermion measurements; new combination when all experiments have final results
- Expect small improvements to $\sigma(q\bar{q})$, but leptons dominated by statistics




WW Cross-sections



- 3 channels: $W^+W^- \rightarrow q\bar{q}q\bar{q}$
 $W^+W^- \rightarrow q\bar{q}l\bar{\nu}_l$
 $W^+W^- \rightarrow l\bar{\nu}_l l\bar{\nu}_l$

- LEP combination updated with final values from **ALEPH, L3**
 Final combination awaiting **OPAL** final results




Good agreement with theoretical expectations

W Branching Ratios

- LEP combination updated with final values from ALEPH, L3


Summer 2004 - LEP Preliminary

W Hadronic Branching Ratio



Summer 2004 - LEP Preliminary


W Leptonic Branching Ratios






- $B(W \rightarrow \tau \nu)$ higher than average of $B(W \rightarrow \mu \nu)$ and $B(W \rightarrow e \nu)$ by 3σ

Single Vector Boson Production

• **Zee**




• **Weν**




LEP combination updated with final measurements from **ALEPH, L3**

Charged Triple Gauge Boson Couplings



- Measured using WW events:
 $\sigma(WW)$, $\cos \theta_W$, W decay angles
- Also $W e \nu$ and $\nu \bar{\nu} \gamma$ channels
- Assuming C, P conservation and gauge constraints: $14 \rightarrow 3$ couplings: $g_1^Z, \lambda_\gamma, \kappa_\gamma$
- LEP combination: combine $\log \mathcal{L}$ curves including correlated systematics




Charged Triple Gauge Boson Couplings

- L3, OPAL values final, ALEPH, DELPHI values preliminary
- LEP combined results allowing one free parameter:

$$\begin{aligned}
 g_1^Z &= 0.991^{+0.022}_{-0.021} \\
 \kappa_\gamma &= 0.984^{+0.042}_{-0.047} \\
 \lambda_\gamma &= -0.016^{+0.021}_{-0.023}
 \end{aligned}$$



- Good agreement with Standard Model
- Couplings measured with precision of 2–4%



- LEP combined confidence levels allowing 2 free parameters

Other Gauge Boson Couplings



- Neutral Triple Gauge Boson couplings ($ZZ\gamma$, $Z\gamma\gamma$) are zero in SM
- Limits set from ZZ and $Z\gamma$ channels
- SM Quartic Gauge Couplings either zero or too small to be observed at LEP
- Limits set from $WW\gamma$, $Z\gamma\gamma$ and $\nu\bar{\nu}\gamma\gamma$ channels



W Mass Measurement

- A principal aim of LEP2
- Comparison of direct measurement with indirect determination from EW fits is test of SM
- Measure by direct reconstruction of $q\bar{q}$ or $l\bar{\nu}_l$ mass in $W^+W^- \rightarrow q\bar{q}l\bar{\nu}_l$ and $W^+W^- \rightarrow q\bar{q}q\bar{q}$ channels
- Reconstruct event-by-event mass using beam energy constraint (kinematic fit) to improve resolution

OPAL 183-209 GeV $\int L dt = 677 \text{ pb}^{-1}$



- Fit mass distribution $\Rightarrow m_W$, using MC to correct for bias


W Mass Measurement

- LEP preliminary values:

$$m_W(q\bar{q}\ell\bar{\nu}_\ell) = 80.411 \pm 0.032(\text{stat}) \pm 0.030(\text{sys}) \text{ GeV}$$

$$m_W(q\bar{q}q\bar{q}) = 80.420 \pm 0.035(\text{stat}) \pm 0.101(\text{sys}) \text{ GeV}$$

Summer 2003 - LEP Preliminary



- Systematics completely dominate $q\bar{q}q\bar{q}$ channel (and important in $q\bar{q}\ell\bar{\nu}_\ell$ channel)
- Combined value:


$$m_W = 80.412 \pm 0.042 \text{ GeV}$$
- What has been happening in last year to improve and finalize measurements?

W Mass Measurement


- Main systematics in $q\bar{q}q\bar{q}$ channel arise from **final state interactions**
- Separation of W decay vertices $\sim 0.1 \text{ fm} <$ hadronic scale $\sim 1 \text{ fm}$
 \Rightarrow W decays have space-time overlap and can exchange colour:

Colour Reconnection


- May also be **Bose-Einstein Correlations** between like-sign particles from different W's
- Both effects may shift measured m_W by large amount ($\sim 100 \text{ MeV}$)
Not included in standard MC used to calibrate m_W measurement
- Much effort to measure CR and BEC effects to estimate realistic errors



Colour Reconnection



- Effects estimated using phenomenological models
- Look at particle flow between jets in $W^+W^- \rightarrow q\bar{q}q\bar{q}$ events
- Compare particle flow in inter-W region with intra-W region



- L3 results final, ADO preliminary


- Use largest reconnection probability compatible with data to set $\Delta(m_W)$
- Current LEP combination uses $\kappa = 2.1$ in SK1 model ($\sim 55\%$ reconnected)

$$\Rightarrow \Delta m_W(q\bar{q}q\bar{q}) = 90 \text{ MeV}$$

Colour Reconnection

- Soft particles most affected by FSI
⇒ reduce effects with momentum cuts or jet cone cuts
Reduces FSI error at expense of statistical error, as jet directions less precisely determined
- Final m_W analyses will optimize jet reconstruction

DELPHI preliminary SK1 curves



Bose-Einstein Correlations

- BEC between like-sign pions well-established in Z decays
Do they occur between particles from different W's?
- Currently contributes 35 MeV to $\Delta m_W(q\bar{q}q\bar{q})$ (LUBOEI model)

Bose-Einstein Correlations



- Study using two-particle correlations, normalized to 'no BEC'
- Use $W^+W^- \rightarrow q\bar{q}l\bar{\nu}_l$ events to estimate intra- W correlations, mixed events (or MC) for kinematic correlations

$$\Delta\rho(Q) = \rho_2^{WW}(Q) - 2\rho_2^W(Q) - 2\rho_{\text{mix}}^{WW}(Q)$$

- LEP combined data:

$$\frac{\text{data} - \text{noBE}}{\text{fullBE} - \text{noBE}} = 0.23 \pm 0.13$$

- DELPHI, L3, OPAL results now final
- L3, OPAL compatible with no inter- W BEC, DELPHI sees significant effect



Prospects for Final m_W Measurement


- LEP measurements of m_W should be finalized soon
- Expect improvements over preliminary measurement
- LEP beam energy uncertainty will decrease from 21 MeV to ~ 10 MeV
Final LEP beam energy paper recently submitted for publication
- Analyses in $q\bar{q}q\bar{q}$ channel will be optimized to reduce total error
- Hadronization and detector effects will be better understood
- In absence of systematics, LEP statistical precision ~ 21 MeV
With reduced weight for $q\bar{q}q\bar{q}$ channel, probably ~ 25 MeV
- Final total error will probably be in range 32–40 MeV, depending on FSI results

W Width Measurement

- Fits to W mass distributions also determine Γ_W
- Preliminary LEP average:

$$\Gamma_W = 2.150 \pm 0.091 \text{ GeV}$$

Summer 2003 - LEP Preliminary



Standard Model Fit

- Standard Model fit by LEPEWWG uses 17 inputs from LEP, SLD, Tevatron:

Z lineshape: $m_Z, \Gamma_Z, \sigma_{\text{had}}^0, R_\ell, A_{\text{fb}}^{0,\ell}$

τ polarisation: P_τ

Polarised lepton asymmetry: $\mathcal{A}_\ell(\text{SLD})$

Heavy flavour: $R_b, R_c, A_{\text{FB}}^{0,b}, A_{\text{FB}}^{0,c}, \mathcal{A}_b, \mathcal{A}_c$

Inclusive hadronic charge asymmetry

m_t, m_W, Γ_W

- Updates since summer 2003:

- Tevatron m_t : $178.0 \pm 2.7 \pm 3.3$ GeV

- LEP1 heavy flavours


- Theory calculations including full two-loop corrections for m_W and $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ (Awramik, Czakon, Freitas, Weiglein)

Shifts predicted value of m_H from $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ alone by ~ 19 GeV



Standard Model Fit

Summer 2004

- Fit $\chi^2/\text{dof} = 15.8/13$
- 67% correlation between m_t and $\log m_H$
- Largest contribution to χ^2 from $A_{\text{FB}}^{0,b}$ (2.4σ)
- $A_{\text{FB}}^{0,b}$ prefers large m_H , whereas R_ℓ , m_W and lepton asymmetries prefer small m_H



Standard Model Fit



Good agreement between measured m_W , m_t and values predicted by fit excluding direct measurements

$m_H < 260$ GeV 95% c.l.

QCD and Two-photon Physics


Will discuss:

- α_s from event shapes
- α_s from 4-jet rate
- Unbiased gluon jets
- Coherence effects in $Z \rightarrow 3$ jets
- Dead cone effect
- Inclusive jet/hadron production in two-photon events

No time to cover

- Fragmentation functions and scaling violation
- Fermi-Dirac and Bose-Einstein correlations in Z decays
- b-quark mass effects
- Production of Ξ_c^0 , Ξ_b in Z decays
- Pentaquark searches


α_s from Event Shapes



- Event shapes sensitive to gluon emission, e.g. thrust:



$$T = \max_{\vec{n}} \left(\frac{\sum_i |p_i \cdot \vec{n}|}{\sum_i |p_i|} \right)$$

- Fit event shape distributions with $\mathcal{O}(\alpha_s^2)$ + NLLA (log R matching) QCD predictions $\Rightarrow \alpha_s$
- Final measurements from all experiments




α_s from Event Shapes

- Combinations by LEPQCD group
- Use variables which are **infra-red safe** (soft gluon emission) and **collinear stable** (collinear parton branchings)
- Combination requires care because of large correlated errors
- Treat hadronization and theory errors as uncorrelated when calculating weights, but include 100% correlation when calculating hadronisation and theory uncertainties on combined α_s




α_s from Event Shapes

- Preliminary combination of final results from all experiments
- Relative weights in global fit:
LEP1 46%
LEP2 54%
- α_s runs as expected from QCD
- Theory error includes variation of renorm. scale: $0.5 < x_\mu < 2$,
log rescaling fact.: $2/3 < x_L < 3/2$,
matching scheme and kinematic cutoffs




$$\alpha_s(m_Z) = 0.1202 \pm 0.0003(\text{stat}) \pm 0.0007(\text{exp}) \pm 0.0015(\text{hadr}) \pm 0.0044(\text{theo})$$



α_s from 4-jet Rate



- New meas. of α_s from 4-jet rate
- **DELPHI**: Cambridge jet algorithm
- Fit to $\mathcal{O}(\alpha_s^3)$ QCD prediction of DEBRECEN (Nagy, Trocsanyi) using experimentally optimized renormalization scale
- **OPAL**: Durham jet finder
- Fit to $\mathcal{O}(\alpha_s^3)$ + NLLA QCD prediction with $x_\mu=1$



α_s from 4-jet Rate



● DELPHI


$$\alpha_s(m_Z) = 0.1175 \pm 0.0030$$

● OPAL


$$\alpha_s(m_Z) = 0.1208 \pm 0.0038$$

Unbiased Gluon Jets with Jet Boost Algorithm


- Jet boost algorithm (Eden, Gustafson) relates gluon jets in $q\bar{q}g$ events to gg system \Rightarrow unbiased gluon jets



(a)



(b)





(c)



- Decompose into 2 colour dipoles
- Boost each dipole into back-to-back frame
- Recombine to give event with gg structure

Unbiased Gluon Jets with Jet Boost Algorithm

- Method used by OPAL to measure gluon charged multiplicity for $5 < E_g < 20$ GeV
- Results consistent with other measurements, and most precise for this energy range
- Theoretical fits OK
- Also measure gluon fragmentation function



Inclusive Charged Particle Distributions




- Mean charged multiplicity
Well-described by theory or MC

- Peak of $\xi = -\ln x_p$ distribution
Sensitive to coherence effects


Coherence Effects in 3-jet Events

- New results from DELPHI
- Measure multiplicity in 30° cone perpendicular to event plane in 3-jet events
- Compare with corresponding quantity in 2-jet events
- Direct evidence for coherence effects in soft particle production




Dead Cone Effect

- QCD predicts gluon radiation off heavy quark suppressed at small angles
- DELPHI study using 2-jet $Z \rightarrow b\bar{b}$ and $Z \rightarrow c\bar{c}$ events
- Remove particles associated with b- or c-quark
- Compare angular distribution of fragmentation particles in b- and c-jets
- First DIRECT observation of 'dead cone' effect



Two-photon Physics


- Active area with many new measurements \Rightarrow test QCD
- e.g. new **DELPHI** measurement of total hadronic cross-section at low Q^2
uses **Very Small Angle Tagger** at $\theta \sim 3-15$ mrad
- Direct event-by-event reconstruction of $W_{\gamma\gamma}$ for double-tagged events without unfolding \Rightarrow **small systematic errors**




- DELPHI measurements somewhat higher than L3 and OPAL

Inclusive jet production

- L3 measurements of inclusive jet production in two-photon collisions for $|\eta| < 1$
 $\langle W_{\gamma\gamma} \rangle \sim 30 \text{ GeV}$
 $\langle Q^2 \rangle \sim 0.2 \text{ GeV}^2$
- p_t spectrum well-represented by power law
- Data higher than NLO QCD prediction (Frixione, Bertora)




Inclusive hadron production



- L3 measurements of inclusive π^\pm production in two-photon collisions for $|\eta| < 1$ and $\langle W_{\gamma\gamma} \rangle > 5 \text{ GeV}$
- Also see excess at high p_t compared with NLO QCD calculations (Binnewies, Kniehl, Kramer)
- Similar disagreement seen in π^0

Inclusive hadron production

- New DELPHI measurement of inclusive hadron production
 $|\eta| < 1.5$
 $5 < W_{\gamma\gamma} < 203 \text{ GeV}$
- Good agreement with NLO QCD for $p_t < 6 \text{ GeV}$
- Some excess of data over NLO QCD at high p_t , but not at L3 level




Exclusive Particle Production

- Several measurements of exclusive particle production
- e.g. **L3** have measured $\rho^0\rho^0$ and $\rho^+\rho^-$
- In range $1.1 < W_{\gamma\gamma} < 2.1$ GeV,
 $1.2 < Q^2 < 8.5$ GeV²

$$\frac{\sigma(\rho^+\rho^-)}{\sigma(\rho^0\rho^0)} = 1.81 \pm 0.47 \pm 0.22$$

expect 2 from isospin

- Fit to QCD expectation $\frac{d\sigma_{ee}}{dQ^2} \sim \frac{1}{Q^n(Q^2 + \langle W_{\gamma\gamma} \rangle^2)^2}$
gives $n = 2.4 \pm 0.3$ (2.5 ± 0.4) for $\rho^0\rho^0$ ($\rho^+\rho^-$) (expect $n = 2$)



Summary

- Since LEP finished running in 2000, the experiments have continued to produce lots of new physics results covering a wide range of topics:
Fermion- and boson-pair cross-sections, gauge boson couplings, QCD, two-photon physics....
- Some important LEP2 measurements still to be finalized, especially m_W
- Important to utilize fully the plentiful, high-quality data produced at LEP