

**$B_J(5970)^+$** 

$$I(J^P) = \frac{1}{2}(??)$$

$I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

 **$B_J(5970)^+$  MASS**OUR FIT uses  $m_{B^0}$  and  $m_{B_J(5970)^+} - m_{B^0}$  to determine  $m_{B_J(5970)^+}$ .

VALUE (MeV)	DOCUMENT ID
<b>5964 ± 5 OUR FIT</b>	

 **$m_{B_J(5970)^+} - m_{B^0}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**685 ± 5 OUR FIT****685 ± 5 OUR AVERAGE**

685.3 ± 4.1 ± 2.5	2K	<sup>1</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
681 ± 5 ± 12	1.4k	<sup>2</sup> AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.8 ± 4.5 ± 2.5	2K	<sup>3</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
-------------------	----	-------------------	-----------	------------------

<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses two relativistic Breit-Wigner functions in the fit for mass difference.<sup>2</sup> AALTONEN 14l reports  $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$  MeV which we adjusted by the  $\pi^+$  mass.<sup>3</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$m_{B_J(5970)^+} - m_{B^{*0}}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.0 ± 4.0 ± 2.5	2k	<sup>1</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
-------------------	----	-------------------	-----------	------------------

<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$  MeV whichwe adjust by the  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = -(-1)^J$ ,  $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

**$B_J(5970)^+$  WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>62 \pm 20</math> OUR AVERAGE</b>				
$63 \pm 15 \pm 17$	2K	<sup>1</sup> AAIJ	15AB LHCB	$p p$ at 7, 8 TeV
$60^{+30}_{-20} \pm 40$	1.4k	AALTONEN	14i CDF	$p \bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$61 \pm 14 \pm 17$	2K	<sup>2</sup> AAIJ	15AB LHCB	$p p$ at 7, 8 TeV
$61 \pm 15 \pm 17$	2K	<sup>3</sup> AAIJ	15AB LHCB	$p p$ at 7, 8 TeV

<sup>1</sup> Assuming  $P = (-1)^J$  and using two relativistic Breit-Wigner functions in the fit for mass difference.

<sup>2</sup> Assuming  $P = (-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

<sup>3</sup> Assuming  $P = -(-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

 **$B_J(5970)^+$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^0 \pi^+$	possibly seen
$\Gamma_2$ $B^{*0} \pi^+$	seen

 **$B_J(5970)^+$  BRANCHING RATIOS**

<u><math>\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}</math></u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_1/\Gamma$
possibly seen	2K	<sup>1</sup> AAIJ	15AB LHCB	$p p$ at 7, 8 TeV	
<b>possibly seen</b>	1.4k	AALTONEN	14i CDF	$p \bar{p}$ at 1.96 TeV	

<sup>1</sup> A  $B\pi$  decay is forbidden from a  $P = -(-1)^J$  parent, whereas  $B^*\pi$  is allowed.

<u><math>\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}</math></u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_2/\Gamma$
seen	2k	AAIJ	15AB LHCB	$p p$ at 7, 8 TeV	
<b>seen</b>	1.4k	AALTONEN	14i CDF	$p \bar{p}$ at 1.96 TeV	

 **$B_J(5970)^+$  REFERENCES**

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14i PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)