

$B_1(5721)^+$

$$I(J^P) = \frac{1}{2}(1^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_1(5721)^+$ MASSOUR FIT uses $m_{B^{*0}}$ and $m_{B_1^+} - m_{B^{*0}}$ to determine $m_{B_1(5721)^+}$.

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
$5725.9^{+2.5}_{-2.7}$ OUR FIT	

 $m_{B_1^+} - m_{B^{*0}}$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

 $401.2^{+2.4}_{-2.7}$ OUR FIT **$401.2^{+2.4}_{-2.7}$ OUR AVERAGE**

$400.5 \pm 1.8 \pm 3.1$	8K	¹ AAIJ	15AB	LHCB	$p\bar{p}$ at 7, 8 TeV
$402 \pm 3 \pm 1_{-3}$		² AALTONEN	14I	CDF	$p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_1^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^0}) - m_{\pi^+} = 260.9 \pm 1.8 \pm 3.1$ MeV which we adjust by the π^+ mass and assume $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.² AALTONEN 14I reports $m_{B_1(5721)^+} - m_{B^{*0}} - m_{\pi^+} = 262 \pm 3^{+1}_{-3}$ MeV which we adjusted by the π^+ mass. **$B_1(5721)^+$ WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

 31 ± 6 OUR AVERAGE Error includes scale factor of 1.1.

$29.1 \pm 3.6 \pm 4.3$	8K	AAIJ	15AB	LHCB	$p\bar{p}$ at 7, 8 TeV
$49 \pm 12 \pm 2_{-10 \ -13}$		AALTONEN	14I	CDF	$p\bar{p}$ at 1.96 TeV

 $B_1(5721)^+$ DECAY MODES

<u>Mode</u>	<u>Fraction (Γ_i/Γ)</u>
$\Gamma_1 \quad B^{*0} \pi^+$	seen

$B_1(5721)^+$ BRANCHING RATIOS

$\Gamma(B^{*0}\pi^+)/\Gamma_{\text{total}}$					Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	8K	AAIJ	15AB LHCb	pp at 7, 8 TeV	
seen		AALTONEN	14i CDF	$p\bar{p}$ at 1.96 TeV	

$B_1(5721)^+$ REFERENCES

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14i	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)