

**$\psi(4660)$**

$I^G(J^{PC}) = 0^-(1^{--})$

also known as  $Y(4660)$ ; was  $X(4660)$

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

Seen in radiative return from  $e^+e^-$  collisions at  $\sqrt{s} = 9.54\text{--}10.58$  GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

### **$\psi(4660)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4633 <math>\pm</math> 7 OUR AVERAGE</b>	Error includes scale factor of 1.4. See the ideogram below.			
4625.9 $\pm$ 6.2	$\pm$ 0.4	89	<sup>1</sup> JIA	19A BELL $e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652	$\pm$ 10	$\pm$ 11	279	<sup>2</sup> WANG
4669	$\pm$ 21	$\pm$ 3	37	<sup>3</sup> LEES
4634	$\pm$ 8	$\pm$ 5	142	<sup>4</sup> PAKHLOVA
	$\pm$ 7	$\pm$ 8		08B BELL $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4652.5 $\pm$ 3.4 $\pm$ 1.1	5	DAI	17	RVUE $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$
4645.2 $\pm$ 9.5 $\pm$ 6.0	6	ZHANG	17B	RVUE $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4646.4 $\pm$ 9.7 $\pm$ 4.8	7	ZHANG	17C	RVUE $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
4661	$\pm$ 9	$\pm$ 6	44	<sup>8</sup> LIU
	$\pm$ 8			08H RVUE $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4664	$\pm$ 11	$\pm$ 5	44	WANG
				07D BELL $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

<sup>1</sup> From a fit of a Breit-Wigner convolved with a Gaussian.

<sup>2</sup> From a two-resonance fit. Supersedes WANG 07D.

<sup>3</sup> From a two-resonance fit.


<sup>4</sup> The  $\pi^+\pi^-\psi(2S)$  and  $\Lambda_c^+\Lambda_c^-$  states are not necessarily the same.

<sup>5</sup> The pole parameters are extracted from the speed plot.

<sup>6</sup> From a three-resonance fit.

<sup>7</sup> From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+\pi^-J/\psi$  and  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  data.

<sup>8</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.



### $\psi(4660)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>64 <math>\pm</math> 9 OUR AVERAGE</b>				
49.8 $^{+13.9}_{-11.5}$ $\pm$ 4.0	89	<sup>1</sup> JIA	19A BELL	$e^+ e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
68 $\pm$ 11 $\pm$ 5	279	<sup>2</sup> WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
104 $\pm$ 48 $\pm$ 10	37	<sup>3</sup> LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
92 $^{+40}_{-24}$ $^{+10}_{-21}$	142	<sup>4</sup> PAKHLOVA	08B BELL	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62.6 $\pm$ 5.6 $\pm$ 4.3		<sup>5</sup> DAI	17 RVUE	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
113.8 $\pm$ 18.1 $\pm$ 3.4		<sup>6</sup> ZHANG	17B RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
103.5 $\pm$ 15.6 $\pm$ 4.0		<sup>7</sup> ZHANG	17C RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
42 $^{+17}_{-12}$ $\pm$ 6	44	<sup>8</sup> LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
48 $\pm$ 15 $\pm$ 3	44	WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

<sup>1</sup> From a fit of a Breit-Wigner convolved with a Gaussian.

<sup>2</sup> From a two-resonance fit. Supersedes WANG 07D.

<sup>3</sup> From a two-resonance fit.

<sup>4</sup> The  $\pi^+ \pi^- \psi(2S)$  and  $\Lambda_c^+ \Lambda_c^-$  states are not necessarily the same.

<sup>5</sup> The pole parameters are extracted from the speed plot.

<sup>6</sup> From a three-resonance fit.

<sup>7</sup> From a combined fit of BELLE, BABAR and BES3  $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$  and  $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$  data.

<sup>8</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

## $\psi(4660)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 e^+ e^-$	not seen
$\Gamma_2 \psi(2S) \pi^+ \pi^-$	seen
$\Gamma_3 J/\psi \eta$	not seen
$\Gamma_4 D^0 D^{*-} \pi^+$	not seen
$\Gamma_5 \chi_{c1} \gamma$	not seen
$\Gamma_6 \chi_{c2} \gamma$	not seen
$\Gamma_7 \Lambda_c^+ \Lambda_c^-$	seen
$\Gamma_8 D_s^+ D_{s1}(2536)^-$	seen

$$\psi(4660) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$$

$$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_2 \Gamma_1 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
2.0 $\pm$ 0.3 $\pm$ 0.2	279	<sup>1</sup> WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
8.1 $\pm$ 1.1 $\pm$ 1.0	279	<sup>2</sup> WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.7 $\pm$ 1.3 $\pm$ 0.5	37	<sup>3</sup> LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.5 $\pm$ 1.7 $\pm$ 0.7	37	<sup>4</sup> LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.2 $\pm$ 0.7 -0.6	44	<sup>5</sup> LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
5.9 $\pm$ 1.6	44	<sup>6</sup> LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
3.0 $\pm$ 0.9 $\pm$ 0.3	44	<sup>3</sup> WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.6 $\pm$ 1.8 $\pm$ 0.8	44	<sup>4</sup> WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

<sup>1</sup> Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

<sup>2</sup> Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

<sup>3</sup> Solution I of two equivalent solutions in a fit using two interfering resonances.

<sup>4</sup> Solution II of two equivalent solutions in a fit using two interfering resonances.

<sup>5</sup> Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

<sup>6</sup> Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_3 \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

**• • •** We do not use the following data for averages, fits, limits, etc. **• • •**

$$<0.94 \quad 90 \quad \text{WANG} \quad 13B \text{ BELL} \quad e^+ e^- \rightarrow J/\psi \eta \gamma$$

$$\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_5 \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$$<\mathbf{0.45} \quad 90 \quad ^1 \text{ HAN} \quad 15 \text{ BELL} \quad 10.58 \text{ } e^+ e^- \rightarrow \chi_{c1} \gamma$$

<sup>1</sup> Using  $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$ .

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_6\Gamma_1/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.1	90	1 HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$
<sup>1</sup> Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .					
$\Gamma(D_s^+ D_{s1}(2536)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_8\Gamma_1/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>14.3^{+2.8}_{-2.6} \pm 1.5</math></b>	89	1 JIA	19A	BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
<sup>1</sup> Using $D_{s1}(2536)^- \rightarrow \bar{D}^{*0} K^-$ .					

## $\psi(4660)$ BRANCHING RATIOS

$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S)\pi^+ \pi^-)$				$\Gamma_4/\Gamma_2$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<10	90	PAKHLOVA	09	BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					
VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_4/\Gamma \times \Gamma_1/\Gamma$	
<b><math>&lt;0.37 \times 10^{-6}</math></b>	90	1 PAKHLOVA	09	BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$
<sup>1</sup> Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$ .					
$\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$	
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>0.68^{+0.16+0.29}_{-0.15-0.30}</math></b>	142	1 PAKHLOVA	08B	BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
<sup>1</sup> The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.					

## $\psi(4660)$ REFERENCES

JIA	19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)
DAI	17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan	
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)