

## **Reversible and irreversible processes**



#### **Carnot engine**






https://courses.lumenlearning.com/boundless-physics/chapter/the-second-law-of-thermodynamics/

# **Carnot engine**

Thermal efficiency of the engine:



$$e = 1 - \frac{|Q_C|}{|Q_H|}$$

An ideal gas is taken through a Carnot cycle. The isothermal expansion occurs at 250°C, and the isothermal compression takes place at 50.0°C. The gas takes in 1.20x10<sup>3</sup> J of energy from the hot reservoir during the isothermal expansion. Find (a) the energy expelled to the cold reservoir in each cycle and (b) the net work done by the gas in each cycle.


In which case will the efficiency of a Carnot cycle be higher: When the hot body temperature is increased by  $\Delta T$ , or when the cold body temperature is decreased by the same magnitude?

# **Carnot's principle**

#### **Carnot's principle**

"The efficiency of a quasi-static or reversible Carnot cycle depends only on the temperatures of the two heat reservoirs, and is the same, whatever the working substance. A Carnot engine operated in this way is the most efficient possible heat engine using those two temperatures."

## **Carnot Engine and the Concept of Entropy**




## **State variables**

Do you understand the following statement?

- Internal energy is a state variable
- Heat is not a state variable
- Entropy is a state variable

You can start by considering the cyclic process



https://physics.stackexchange.com/questions/370468

N. Srimanobhas; Heat engines, entropy, and the second law of thermodynamics

Imagine a Carnot engine that operates between the temperatures  $T_H = 850$  K and  $T_L = 300$  K. The engine performs 1200 J of work each cycle, which takes 0.25 s. (a) What is the efficiency of this engine?

(b) What is the average power of this engine?

- (c) How much energy  $IQ_HI$  is extracted as heat from the high-temperature reservoir every cycle
- (d) How much energy IQ<sub>L</sub>I is delivered as heat to the low- temperature reservoir every cycle?
- (e) By how much does the entropy of the working substance change as a result of the energy transferred to it from the high-temperature reservoir? From it to the low-temperature reservoir?

## Entropy

## **2nd Law of the Thermodynamics**

The second law of thermodynamics states that a closed system has entropy that may

increase: irreversible process

or otherwise

remain constant: reversible process

lf

$$\Delta S = \Delta S_{\text{gas}} + \Delta S_{\text{res}}$$

And processes can happen if

 $\Delta S \ge 0$ 

When a metal bar is connected between a hot reservoir at  $T_h$  and a cold reservoir at  $T_c$ , the energy transferred by heat from the hot reservoir to the cold reservoir is Q. In this irreversible process, find expressions for the change in entropy of (a) the hot reservoir, (b) the cold reservoir, and (c) the Universe, neglecting any change in entropy of the metal rod.

## **Difference between reversible/irreversible**

A weightless piston divides a thermally insulated cylinder into two equal parts. One part contains one mole of an ideal gas, the other is evacuated. The piston is released and the gas fills the whole volume of the cylinder. Then the piston is slowly displaced back to the initial position. Find the increment of the entropy of the gas resulting from these two processes.